



CP asymmetry in $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ decays

T. N. Pham^a

^a Centre de Physique Theorique,
Centre National de la Recherche Scientifique, UMR 7644,
Ecole Polytechnique, 91128 Palaiseau Cedex, France

The near equality between the measured value for $|V_{ub}|/|V_{cb}|$ and its unitarity triangle lower bound from $\sin 2\beta$ measurements suggests that $\alpha \approx 90^\circ$ and $\sin \gamma \approx \cos \beta = 0.91$ and a large direct CP violation in B decays would be possible. In this talk, I would like to discuss a recent analysis of the non resonant and resonant CP asymmetry for the decays $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ where the CP asymmetry at the χ_{c0} resonance comes from the interference of the non resonant amplitude with the resonant amplitude $B^\pm \rightarrow \chi_{c0} K^\pm \rightarrow \pi^+ \pi^- K^\pm$ and $B^\pm \rightarrow \chi_{c0} K^\pm \rightarrow K^+ K^- K^\pm$. Using the the Babar and Belle measurements of the $B^- \rightarrow \chi_{c0} K^-$ branching ratios, we predict that partial width asymmetry at the χ_{c0} resonance could reach 10% for the $B^- \rightarrow \pi^+ \pi^- K^-$ decay and 16% for the $B^- \rightarrow K^+ K^- K^-$ decay.

1 Introduction

In three-body charmless B decays, e.g $B \rightarrow K\pi\pi$, apart from the non resonant CP asymmetry there is also the partial width CP asymmetries produced by the interference between the non resonant and resonant amplitudes, both of them could be tree amplitudes. Previous studies [1, 2, 3] show that in $B^- \rightarrow \pi^+ \pi^- \pi^-$ decays, a large partial width CP asymmetry at the χ_{c0} resonance could arise from the interference of the non resonant amplitude with the resonant amplitude coming from the decays $B^- \rightarrow \chi_{c0} \pi^- \rightarrow \chi_{c0} \rightarrow \pi^- \pi^-$, but no firm prediction for the asymmetry could be given as there is no theoretical prediction for the $B^- \rightarrow \chi_{c0} \pi^-$ decay rate. A similar analysis can now be made for the CP asymmetries in the $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ decays using the recent Belle[4] and Babar[5] measurements of the $B^- \rightarrow \chi_{c0} K^-$ branching ratios: $\text{BR}(B^- \rightarrow \chi_{c0} K^-) = (6.0^{+2.1}_{-1.8}) \times 10^{-4}$ $\text{BR}(B^- \rightarrow \chi_{c0} K^-) = (2.4 \pm 0.7) \times 10^{-4}$ for Belle and Babar respectively. The non resonant branching ratios reported by Belle[6] are: $\text{BR}(B^- \rightarrow \pi^+ \pi^- K^-) = (58.5 \pm 7.1 \pm 8.8) \times 10^{-6}$ $\text{BR}(B^- \rightarrow K^+ K^- K^-) = (37.0 \pm 3.9 \pm 4.4) \times 10^{-6}$. In this talk, I would like to discuss a recent work[7] on the non resonant CP asymmetries and the partial CP asymmetries at the χ_{c0} resonance in these decays.

2 The CKM matrix

The effective weak Hamiltonian for the nonleptonic Cabibbo-suppressed B meson decays is given by[8, 9, 10]

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} [V_{ub} V_{us}^* (c_1 O_{1u} + c_2 O_{2u}) + V_{cb} V_{cs}^* (c_1 O_{1c} + c_2 O_{2c})] - \sum_{i=3}^{10} ([V_{ub} V_{us}^* c_i^u + V_{cb} V_{cs}^* c_i^c + V_{tb} V_{ts}^* c_i^t] O_i) + \text{h.c.} \quad (1)$$

where O_1, O_2 are the tree-level operators and $O_3 - O_6$ are the penguin operators. As usual with the factorization model we use in our analysis, the hadronic matrix elements are obtained from the effective Hamiltonian in Eq.(1) with c_i replaced by a_i where c_i are next-to-leading Wilson coefficients. Since a_3 and a_5 are one order of magnitude smaller than a_4 and a_6 , the contributions from O_3 and O_5 can be safely neglected. For $N_c = 3$, $m_b = 5 \text{ GeV}$, we use [9, 10] : $a_1 = 1.05$, $a_2 = 0.07$, $a_4 = -0.043 - 0.016i$, $a_6 = -0.054 - 0.016i$. Before writing down the decay amplitudes, I would like to discuss the current determination of the CKM parameters and show that the unitarity triangle lower bound for $|V_{ub}|/|V_{cb}|$ obtained from the Babar and Belle measurements of $\sin 2\beta$ and the measured value for $|V_{ub}|/|V_{cb}|$ already tells us that $\alpha \approx 90^\circ$ and $\sin \gamma \approx \cos \beta = 0.91$. The CKM parameters in Eq.(1) are the flavor-changing charged current couplings of the weak gauge boson $W^\pm$ with the quarks as given by the Cabibbo-Kobayashi-Maskawa (CKM) unitary quark mixing matrix relates the the weak interaction eigenstate d', s', b' of the charge $Q = -1/3$ quarks to their mass eigenstate d, s, b [11]:

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} \quad (2)$$

The unitarity relation

$$V_{ud} V_{ub}^* + V_{cd} V_{cb}^* + V_{td} V_{tb}^* = 0 \quad (3)$$

can be written as

$$R_b e^{i\gamma} + R_t e^{-i\beta} = 1 \quad (4)$$

and is represented by the (db) unitarity triangle shown in Fig.1. $V_{ub} = |V_{ub}| e^{-i\gamma}$, $V_{td} = |V_{td}| e^{-i\beta}$, and $R_b = |V_{ud} V_{ub}^*|/|V_{cd} V_{cb}^*|$ and $R_t = |V_{td} V_{tb}^*|/|V_{cd} V_{cb}^*|$ [12]

As can be seen from Fig.1,

$$R_b \sin \alpha = \sin \beta, \quad R_t \sin \alpha = \sin \gamma \quad (5)$$

Hence the following lower limits for R_b and R_t :

$$R_b \geq \sin \beta, \quad R_t \geq \sin \gamma \quad (6)$$

This gives:

$$\begin{aligned} |V_{ud}V_{ub}^*| &\geq |V_{cd}V_{cb}^*| \sin \beta \\ |V_{td}V_{tb}^*| &\geq |V_{cd}V_{cb}^*| \sin \gamma \end{aligned} \quad (7)$$

The current measured value of $\sin 2\beta$ gives[13]

$$\sin 2\beta = 0.734 \pm 0.06 \quad (8)$$

from which we obtain two solutions for $\sin \beta$:

$$\sin \beta = \begin{cases} 0.40 \pm 0.04 \\ 0.91 \pm 0.02 \end{cases} \quad (9)$$

On the other hand, $|V_{ub}| = (4.11 \pm 0.25 \pm 0.78) \times 10^{-3}$ from LEP and CLEO inclusive data which is above the value $(3.25 \pm 0.32 \pm 0.64) \times 10^{-3}$ from the CLEO exclusive data[11]. With the average value $|V_{ub}| = (3.60 \pm 0.70) \times 10^{-3}$ from the inclusive and exclusive measurements[11], we find

$$R_b = (0.379 \pm 0.07) \quad (10)$$

which is consistent with the lower limit of $\sin \beta = 0.40 \pm 0.04$. This also excludes the solution $\sin \beta = 0.91$. Conversely, from the measured value for $|V_{cb}|$, a lower limit for $|V_{ub}|$ from $\sin \beta$ is then

$$|V_{ub}| = (3.8 \pm 0.6) \times 10^{-3} \quad (11)$$

Note that the inclusive data is consistent with the lower bound for $|V_{ub}|$ from $\sin 2\beta$ measurement while the exclusive data is slightly below the lower bound. Thus to within




Figure 1. The (db) Unitarity Triangle

experimental errors, the average value for $|V_{ub}|$ suggests that $R_b \approx \sin \beta$. This implies

$$\sin \alpha \approx 1, \quad \alpha \approx \pi/2, \quad R_t \approx \cos \beta \quad (12)$$

This is supported by a recent analysis[12] which gives: $\sin 2\alpha = 0.05 \pm 0.31$, $R_b = 0.404 \pm 0.023$, $R_t = 0.927 \pm 0.061$ which indeed shows that $\alpha \approx \pi/2$ and $R_b \approx \sin \beta$. If α is known, Eq.(5) can be used to extract R_b and R_t . We find

$$\begin{aligned} R_b &= \sin \beta \sqrt{1 + (\cos \alpha / \sin \alpha)^2}, \\ R_t &= \sin \gamma \sqrt{1 + (\cos \alpha / \sin \alpha)^2} \end{aligned} \quad (13)$$

This provides a good determination of $|V_{ub}|$ and $|V_{td}|$ in terms of $\sin \beta$ and $\sin \gamma$, respectively. since $(\cos \alpha)^2$ is of the order a few percent. With $\alpha = 90^\circ$, $\sin \gamma = \cos \beta = 0.91 \pm 0.02$, $R_t = 0.91 \pm 0.02$. Thus R_t is also rather well known. for example, the current determination of R_t from ΔM_d [11] gives $|V_{td}V_{tb}^*| = 0.0079 \pm 0.0015$, $R_t = 0.86 \pm 0.25$ which is consistent with the above value for R_t from $\sin 2\beta$ measurements. Since $\sin \gamma$ is close to 1, we expect large direct CP violation in B decays. We now discuss the CP asymmetry in B decays to 3 pseudo-scalar mesons.

3 $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ decays

Consider now the $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ decays. The non resonant amplitude is mainly given by the O_1 , O_4 , O_6 matrix elements and is dominated by the penguin operators as in $B \rightarrow K\pi$ decays, because of the large CKM factors relative to the tree amplitude (Fig.2). With

$$\begin{aligned} O_1 &= (\bar{s}u)_L (\bar{u}b)_L, \quad O_4 = \sum_q (\bar{s}q)_L (\bar{q}b)_L \\ O_6 &= -2 \sum_q (\bar{s}_L q_R) (\bar{q}_R b_L) \end{aligned} \quad (14)$$



Figure 2. Graphs representing the nonleptonic interactions

In the factorization model, the matrix elements of O_1 , O_4 , O_6 can be computed in terms of the $B \rightarrow \pi\pi l\nu$ form factors

given as:

$$\begin{aligned} & \langle \pi^-(p_1) \pi^+(p_2) | (\bar{u}b)_{V-A} | B^-(p_B) \rangle = \mu \\ & = i r (p_B - p_2 - p_1)_\mu + i w_+ (p_2 + p_1)_\mu \\ & + i w_- (p_2 - p_1)_\mu - 2h \varepsilon_{\mu\alpha\beta\gamma} p_B^\alpha p_2^\beta p_1^\gamma. \end{aligned} \quad (15)$$

The B^* pole terms in the form factors w_+ and w_- are ob-




Figure 3. Non resonant and resonant $B \rightarrow K\pi\pi$ decay

tained using the Hybrid model (HQET with the full propagators for the heavy meson pole terms)[14, 15]

$$\begin{aligned} w_+^{nr}(s, t) = & -\frac{g}{f_1 f_2} \frac{f_{B^*} m_{B^*}^{3/2} m_B^{1/2}}{(t - m_{B^*}^2)} \left[1 - \frac{(m_B^2 - m_1^2 - t)}{2m_{B^*}^2} \right] \\ & + \frac{f_B}{2f_1 f_2} - \frac{\sqrt{m_B} \alpha_2}{f_1 f_2} \frac{1}{2m_B^2} (2t + s - m_B^2 - m_3^2 - 2m_1^2), \end{aligned}$$

$$\begin{aligned} w_-^{nr}(s, t) = & \frac{g}{f_1 f_2} \frac{f_{B^*} m_{B^*}^{3/2} m_B^{1/2}}{t - m_{B^*}^2} \left[1 + \frac{(m_B^2 - m_1^2 - t)}{2m_{B^*}^2} \right] \\ & + \frac{\sqrt{m_B} \alpha_1}{f_1 f_2}. \end{aligned}$$

We thus have in the factorization model :

$$\begin{aligned} \langle O_1 \rangle_{nr} = & \langle K^-(p_3) \pi^+(p_1) \pi^-(p_2) | O_1 | B(p_B) \rangle_{nr} \\ = & -[f_3 m_3^2 r^{nr} + \frac{1}{2} f_3 (m_B^2 - m_3^2 - s) w_+^{nr} \\ & + \frac{1}{2} f_3 (s + 2t - m_B^2 - 2m_1^2 - m_3^2) w_-^{nr}] \end{aligned} \quad (16)$$

$\langle O_4 \rangle_{nr}$ is almost the same as $\langle O_1 \rangle_{nr}$. $\langle O_6 \rangle_{nr}$ is given by

$$\begin{aligned} \langle O_6 \rangle_{nr} = & \langle K^-(p_3) \pi^+(p_1) \pi^-(p_2) | O_6 | B(p_B) \rangle_{nr} \\ = & -\left(\frac{\mathcal{B}}{m_B}\right) \left[2 \frac{f_1 f_2}{f_3} m_3^2 r^{nr} + \frac{f_1 f_2}{f_3} (m_B^2 + m_3^2 - s) w_+^{nr} \right. \\ & \left. + \frac{f_1 f_2}{f_3} (s + 2t - m_B^2 - 2m_1^2 - m_3^2) w_-^{nr} \right]. \end{aligned} \quad (17)$$

Thus $\langle O_6 \rangle_{nr} = (2\mathcal{B}/m_B) \langle O_1 \rangle_{nr}$ to a good approximation. The non resonant $B^- \rightarrow K^- \pi^+ \pi^-$ amplitude is then

$$\begin{aligned} \mathcal{M}_{nr} = & \frac{G}{\sqrt{2}} [V_{ub} V_{us}^* a_1 \langle O_1 \rangle_{nr} \\ & - V_{tb} V_{ts}^* (a_4 \langle O_4 \rangle_{nr} + a_6 \langle O_6 \rangle_{nr})]. \end{aligned} \quad (18)$$

In the high di-pion invariant mass region, for example at the χ_{c0} mass, the two pions in an S -wave 0^{++} can also be produced from the decay $B^- \rightarrow \chi_{c0} K^-$ followed by the decay $\chi_{c0} \rightarrow \pi^+ \pi^-$. This will interfere with the non resonant amplitude and produce a CP asymmetry at the χ_{c0} mass. The large $B^- \rightarrow \chi_{c0} K^-$ branching ratio measured by Belle and Babar provides a possibility to look for CP violation in $B \rightarrow K\pi\pi$ decays. This large BR also indicates that nonfactorizable terms are present in B decays[16]. The resonant amplitude is then

$$\begin{aligned} \mathcal{M}_r(B^- \rightarrow \chi_{c0} K^- \rightarrow \pi^+ \pi^- K^-) = \\ \mathcal{M}(B^- \rightarrow \chi_{c0} K^-) \frac{1}{s - m_{\chi_{c0}}^2 + i\Gamma_{\chi_{c0}} m_{\chi_{c0}}} \mathcal{M}(\chi_{c0} \rightarrow \pi^+ \pi^-) \end{aligned} \quad (19)$$

Using the Belle measured $B^- \rightarrow \chi_{c0} K^-$ branching ratio[4], we computed the differential decay rates and CP asymmetries for the two-pion and two-kaon system in the χ_{c0} mass region shown in Figs.1 and Figs.2 (taken from the published work[7]). For the integrated branching ratios we obtain:

$$\text{BR}(B^- \rightarrow K^- \pi^+ \pi^-)_{nr} = T + P + I_1 \cos \gamma + I_2 \sin \gamma. \quad (20)$$

with $T = 7.0 \times 10^{-6}$, $P = 7.5 \times 10^{-5}$, $I_1 = -4.3 \times 10^{-5}$, $I_2 = -1.5 \times 10^{-5}$, For $\sin \gamma = 0.9$, $\cos \gamma = 0.4$, $g = 0.23$, we obtain a branching ratio $\text{BR}(B^- \rightarrow K^- \pi^+ \pi^-)_{nr} = 5.1 \times 10^{-5}$ for the non resonant $B^- \rightarrow K^- \pi^+ \pi^-$ decay. The effective $DD^* \pi$ coupling g used here is smaller than the soft-pion value of 0.57 because of off-shell effects. g and other parameters α_1 and α_2 are chosen to fit the $B \rightarrow \pi$ and $B \rightarrow \rho$ form factors. The above BR is consistent with the experimental upper limits for the non resonant BR. Similarly,

$$\text{BR}(B^- \rightarrow K^- K^+ K^-)_{nr} = T + P + I_1 \cos \gamma + I_2 \sin \gamma. \quad (21)$$




Figure 4. Differential branching ratios for $B^- \rightarrow \pi^+ \pi^- K^-$ vs. s . The curves (a), (b), (c) are $d(B)/ds$, $d(B)/ds + d(\bar{B})/ds$, $d(B)/ds - d(\bar{B})/ds$ against the two-pion invariant mass squared s , for the non resonant, CP symmetric and CP antisymmetric differential branching ratios respectively.

and $T = 3.4 \times 10^{-6}$, $P = 3.7 \times 10^{-5}$, $I_1 = -2.1 \times 10^{-5}$, $I_2 = -7.4 \times 10^{-6}$. This gives $\text{BR}(B^- \rightarrow K^- K^+ K^-)_{nr} = 2.53 \times 10^{-5}$ for the non resonant $B^- \rightarrow K^- K^+ K^-$ decay. The CP asymmetry of the total decay rate is given by

$$A = \frac{\sin \gamma N_1}{N_2 + \cos \gamma N_3}, \quad (22)$$

where $N_1 = -3.0 \times 10^{-5}$, $N_2 = 16.4 \times 10^{-5}$, $N_3 = -8.6 \times 10^{-5}$ for $B^- \rightarrow \pi^+ \pi^- K^-$ and $N_1 = -1.5 \times 10^{-5}$, $N_2 = 8.2 \times 10^{-5}$, $N_3 = -4.2 \times 10^{-5}$ for $B^- \rightarrow K^+ K^- K^-$. and the total integrated asymmetry is then

$$\begin{aligned} A(B^- \rightarrow \pi^+ \pi^- K^-) &= -0.20 \\ A(B^- \rightarrow K^+ K^- K^-) &= -0.20 \end{aligned} \quad (23)$$

Since $\langle O_4 \rangle_{nr} = \langle O_1 \rangle_{nr}$ and $\langle O_6 \rangle_{nr}$ is proportional to $\langle O_1 \rangle_{nr}$ the above results for the non resonant CP asymmetry are essentially independent of the form factors w_+ and w_- . For the partial width CP asymmetry integrated over the χ_{c0} mass region, we find

$$\begin{aligned} A_p(B^\pm \rightarrow K^\pm \pi^+ \pi^-) &= 7.9 \sin \gamma / (73 - 1.2 \cos \gamma) \\ A_p(B^\pm \rightarrow K^\pm K^+ K^-) &= 7.2 \sin \gamma / (41 - 5.6 \cos \gamma) \end{aligned} \quad (24)$$




Figure 5. Differential branching ratios for $B^- \rightarrow K^+ K^- K^-$ vs. s . The curves (a), (b), (c) are $d(B)(NR)/ds$, $d(B)/ds + d(B)/ds$, $d(B)/ds - d(B)/ds$ against the two-kaon invariant mass squared s , for the non resonant, CP symmetric and CP antisymmetric differential branching ratios respectively.

References

1. G. Eilam, M. Gronau and R. R. Mendel, Phys. Rev. Lett. **74** (1995) 4984.
2. N. G. Deshpande, G. Eilam, X. G. He and J. Trampetić, Phys. Rev. D **52** (1995) 5354.
3. B. Bajc, S. Fajfer, R. J. Oakes, T. N. Pham and S. Prelovšek, Phys. Lett. **B** 447 (1999) 313.
4. K. Abe *et al.*, Phys. Rev. Lett. **88** (2002) 031802.
5. B. Aubert *et al.*, hep-ex/0207066.
6. K. Abe *et al.*, hep-ex/0201007.
7. S. Fajfer, R. J. Oakes and T. N. Pham, Phys. Lett. **B** 539 (2002) 67.
8. A. Ali, G. Kramer, and Cai-Dian Lü, Phys. Rev. D **59** (1999) 014005.
9. N. G. Deshpande and X. G. He, Phys. Lett. **B** 336 (1994) 471; N. G. Deshpande, X. G. He, W. S. Hou, and S. Pakvasa, Phys. Rev. Lett. **82** (1999) 2240.
10. C. Isola and T. N. Pham, Phys. Rev. D **62** (2000) 094002.
11. For a review of the CKM model, see, e.g., Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66** (2002) 010001-113.
12. A. Buras, F. Parodi and A. Stocchi, hep-ph/0207101.
13. Y. Nir, Proceedings of the 31st International Conference on High Energy Physics, Amsterdam, Netherland, 24-31 July 2002, hep-ph/0208080. The average value given in this paper is obtained from the Babar and Belle data as well as previous results from ALEPH and OPAL.
14. B. Bajc, S. Fajfer, R. J. Oakes, and T. N. Pham, Phys. Rev. D **58** (1998) 054009.
15. B. Bajc, S. Fajfer, and R. J. Oakes, Phys. Rev. D **53** (1996) 4957; B. Bajc, S. Fajfer, R. J. Oakes, and S. Prelovšek, Phys. Rev. D **56** (1997) 7027.
16. P. Colangelo, F. de Fazio and T. N. Pham, Phys. Lett. **B** 542 (2002) 71.

I would like to thank the Organisers of the CKM03 Workshop for the warm hospitality at Durham and the EURIDICE Collaboration for a grant.